

## **MASSIVE ROTATOR CUFF REPAIR**

### Rehab Protocol

#### Weeks 1-6:

- Sling Immobilization
- Active ROM Elbow, Wrist and Hand
- True Passive (ONLY) ROM Shoulder
  - Supine Elevation in Scapular plane = 140 degrees
  - External Rotation = 40 degrees
- Scapular Stabilization exercises (sidelying)
- Deltoid isometrics in neutral (submaximal) as ROM improves
- No Pulley/Canes until 6 weeks post-op

#### Weeks 6-12:

- Discontinue Sling
- Active Assist to Active ROM Shoulder As Tolerated
  - Elevation in scapular plane and external rotation to tolerance
  - Begin internal rotation as tolerated
  - Light stretching at end ranges
- Cuff Isometrics with the arm at the side
- Upper Body Ergometer

#### Months 3-12:

- Advance to full ROM as tolerated with passive stretching at end ranges
- Advance strengthening as tolerated: isometrics → bands → light weights (1-5 lbs); 8-12 reps/2-3 sets per rotator cuff, deltoid, and scapular stabilizers
- Only do strengthening 3x/week to avoid rotator cuff tendonitis
- Begin eccentrically resisted motions, plyometrics (ex. Weighted ball toss), proprioception (es. body blade)
- Begin sports related rehab at 4 ½ months, including advanced conditioning
- Return to throwing at 6 months
- Throw from pitcher's mound at 9 months
- Collision sports at 9 months
- MMI is usually at 12 months post-op